

SOME REMARKS ON THE CLASS OF BERNSTEIN FUNCTIONS AND SOME SUB-CLASSES

SONIA FOURATI, WISSEM JEDIDI

1. INTRODUCTION

In this paper we summarize some properties that we think complementary to the book recently published of Schilling-Song-Vondracek [4].

We recall the definition of **Bernstein functions** : a function ϕ defined on $[0, +\infty[$ is called a Bernstein function ($\phi \in \mathcal{B}$) if it has the representation

$$\phi(\lambda) = q + d\lambda + \int_{(0, \infty)} (1 - e^{-\lambda x}) \pi(dx), \quad \lambda \geq 0, \quad (1.1)$$

q, d are non negative. The measure π is called the Lévy measure of ϕ . It is a measure supported by $]0, \infty[$ that integrates the function $\min(x, 1)$.

Bernstein functions are more likely called by probabilists "Laplace exponents of infinite divisible non negative sub-distributions" (or "Laplace exponents of subordinators") and the previous representation is their Lévy-Khintchine expression.

2. A UNIFIED VIEW ON SUBCLASSES OF BERNSTEIN FUNCTIONS

In the sequel all the measures will be understood on the space $]0, \infty[$ and their densities, if they have one, are according to the Lebesgue measure on $]0, \infty[$ which will be denoted by dx .

We recall the definition of the Mellin convolution of two measures ν and τ on the space $]0, \infty[$.

$$\nu \circledast \tau(A) = \int_{]0, \infty[^2} 1_A(xy) \nu(dx) \tau(dy), \quad A \text{ borelian of }]0, \infty[.$$

Notice that this integral may be infinite when ν and/or τ are not finite measures.

Theorem 2.1. *Let π be a Lévy measure.*

1) *The measure π has a non increasing density if and only if π is of the form*

$$\pi = 1_{]0, 1]}(x) dx \circledast \nu,$$

where ν is some a Lévy measure.

Date: December 6, 2011.

MSC 2000 subject classifications. 60E07, 60E15.

Key words. Bernstein functions, complete Bernstein functions, Subordinators, Infinite divisible distributions, Generalized Gamma convolution, Hyperbolically completely monotone, Positive stable density....

This work was partially supported by Agence Nationale de la Recherche grant ANR-09-BLAN-0084-01.

2) The measure $x\pi(dx)$ has a non increasing density if and only if π is of the form

$$\pi = 1_{]0,1]}(x) \frac{dx}{x} \otimes \nu,$$

where ν is a measure which integrates the function $f_0(x) = x1_{x \in]0,1]} + \log x 1_{x > 1}$ (in particular ν is a Lévy measure).

3) The measure π has a density of the form $x^{a-1}k(x)$ with $a \in]-1, \infty[$ and k a completely monotonic function such that $\lim_{+\infty} k(x) = 0$ if and only if π has the expression

$$\pi = x^{a-1} e^{-x} dx \otimes \nu,$$

where ν is a measure which integrates the function f_a given by

$$f_a(x) := \begin{cases} x1_{x \in]0,1]} + x^{-a}1_{x > 1} & \text{if } a \in]-1, 0[, \\ x1_{x \in]0,1]} + \log x 1_{x > 1} & \text{if } a = 0, \\ x1_{x \in]0,1]} + 1_{x > 1} & \text{if } a \in]0, \infty[. \end{cases} \quad (2.1)$$

Consequently, ν is a Lévy measure in all cases. Moreover ν is any Lévy measure in case 1) and in case 3) with $a > 0$.

Proof. Notice that if μ has a density h , then $\mu \otimes \nu$ has the density, denoted by $h \otimes \nu$ and valued in $[0, +\infty]$:

$$h \otimes \nu(x) = \int_{]0, \infty[} \frac{1}{y} h\left(\frac{x}{y}\right) \nu(dy), \quad x > 0.$$

1) Using the last expression for $h(x) = u_0(x) = 1_{]0,1]}(x)$, we have

$$u \otimes \nu(x) = \int_x^{+\infty} \frac{\nu(dy)}{y}.$$

Notice that any non increasing function (valued in $[0, +\infty]$) is of the form $u_0 \otimes \nu$ and conversely. Since

$$\int_0^\infty (x \wedge 1) u_0 \otimes \nu(x) = \int_0^1 \frac{x^2}{2} \frac{1}{x} \nu(dx) + \frac{1}{2} \int_1^\infty \frac{1}{y} \nu(dy) + \int_1^\infty \frac{x-1}{x} \nu(dx),$$

we deduce that the measure with density $\int_x^{+\infty} \frac{\nu(dy)}{y}$ is a Lévy measure if and only if ν integrates the function $x \wedge 1$ or, in other words, ν is a Lévy measure.

2) Using the expression of $h \otimes \nu$ with $h(x) = u_1(x) = \frac{1}{x} 1_{]0,1]}(x)$, we have :

$$u_1 \otimes \nu(x) = \frac{1}{x} \int_x^\infty \nu(dy) = \frac{\nu(]x, +\infty[)}{x}.$$

Notice that any function π , valued in $[0, +\infty]$, such that $x\pi(x)$ is non increasing is of the form $u_1 \otimes \nu$ and conversely. After that, note that

$$\int_0^\infty (x \wedge 1) u_1 \otimes \nu(x) dx = \int_0^1 x \nu(dx) + \nu(]1, +\infty[) + \int_1^{+\infty} \log x \nu(dx).$$

Thus, the measure with density $u_1 \otimes \nu(x)$ is a Lévy measure if and only if $\nu(dx)$ integrates $f_0(x) = x1_{x < 1} + \log x 1_{x > 1}$.

3) Let $a \in]-1, \infty[$, k a complete monotone function such that $k(+\infty) = 0$ and $\pi(dx) = x^{a-1} k(x) 1_{]0, \infty[} dx$. By Bernstein theorem, k is the Laplace transform of some measure μ on $[0, \infty[$, $k(x) = \int_{[0, \infty[} e^{-xt} \mu(dt)$. Since $k(+\infty) = \mu(\{0\})$, the measure μ does not charge 0. Then, we can build the measure $\Delta(du) = \mu(du)u^{-a}$ and write

$$k(x) := \int_{]0, \infty[} e^{-xu} u^a \Delta(du), \quad x > 0. \quad (2.2)$$

Let the function

$$h_a(u) := \int_0^\infty (x \wedge u) x^{a-1} e^{-x} dx, \quad u > 0.$$

By Fubini theorem, we have

$$\int_{]0, \infty[} (x \wedge 1) \pi(dx) = \int_{]0, \infty[} (x \wedge 1) x^{a-1} k(x) dx = \int_{]0, \infty[} h_a(u) \frac{\Delta(du)}{u}.$$

We will find the necessary and sufficient conditions on Δ so that the last integral is finite.

First, notice that $h_a(u) \nearrow \Gamma(a+1)$ when $u \rightarrow +\infty$ and then h_a is bounded for any $a > -1$. Second, elementary computations give the following behavior of h_a at the neighborhood of 0 ,

$$\begin{aligned} \lim_{0+} \frac{h_a(u)}{u} &= \Gamma(a) && \text{if } a > 0; \\ 0 < \liminf_{0+} \frac{h_a(u) - u}{u |\log u|} &\leq \limsup_{0+} \frac{h_a(u) - u}{u |\log u|} < \infty, && \text{if } a = 0; \\ \lim_{0+} \frac{h_a(u)}{u^{1-|a|}} &= \frac{1}{|a|} + \frac{1}{1-|a|}, && \text{if } a \in]-1, 0[. \end{aligned}$$

and then π is a Lévy measure iff $\int_1^{+\infty} \frac{\Delta(du)}{u} du < +\infty$ and

$$\Delta([0, 1]) < +\infty, \text{ if } a > 0, \quad \int_{(0, 1]} |\log u| \Delta(du) < \infty, \text{ if } a = 0, \quad \int_{(0, 1]} \frac{\Delta(du)}{u^{|a|}} < \infty, \text{ if } -1 < a < 0.$$

Notice that in each case $\Delta([0, 1]) < +\infty$ and then $\frac{\Delta(du)}{u}$ is a Lévy measure. Define the measure ν as the image of $\Delta(du)$ induced by the function $u \mapsto 1/u$, it is also a Lévy measure. Notice that properties of the measure Δ are equivalent to the fact that the function f_a in the statement of the theorem is integrable for ν . In order to conclude, write

$$x^{a-1} k(x) = x^{a-1} \int_{]0, \infty[} e^{-xu} u^a \Delta(du) = \int_{]0, \infty[} \left(\frac{x}{y}\right)^{a-1} e^{-\frac{x}{y}} \frac{\nu(dy)}{y} = [y^{a-1} e^{-y} \otimes \nu](x).$$

□

Remark 2.2. Bernstein functions ϕ associated by (1.1) to Lévy measures π of theorem 2.1, are particular in the context of subordinators and infinitely divisible random variables :

1) A Bernstein function whose Lévy measure is of type (1) is called Jurek Bernstein functions. The Jurek class is also characterized by $\phi \geq 0$ and $(\lambda\phi)' \in \mathcal{B}$.

2) A Bernstein function whose Lévy measure is of type (2) is called self-decomposable Bernstein functions, we denote \mathcal{SDB} their set. It is easy to check (see theorem 2.6 ch. VI [5], for instance), that

$$\phi \in \mathcal{SDB} \iff \phi(0) \geq 0 \quad \text{and} \quad \lambda \mapsto \lambda\phi'(\lambda) \in \mathcal{B}.$$

The infinite divisible positive variables X associated to the \mathcal{SDB} -functions ϕ ($\mathbb{E}(e^{-\lambda X}) = e^{-\phi(\lambda)}$) are the self-decomposable r.v. X , that is the random variable for which there exists a family of positive r.v. $(Y_c)_{0 < c < 1}$, each Y_c is independent from X and satisfy $X \stackrel{d}{=} cX + Y_c$.

3) In the book of Schilling-Song-Vondracek the set \mathcal{CB} ([4] p. 49) of Complete Bernstein functions corresponds to the Bernstein functions appearing in part (3) of the above theorem with parameter $a = 1$. The \mathcal{TB} ([4] p. 73) of Thorin Bernstein functions are the ones that with parameter $a = 0$. We will talk again in the next section about CB functions. The class \mathcal{TB} corresponds to the Laplace exponents of the generalized Gamma distributions, shortly GGC, introduced by Bondesson [1], [2] and the GGC subordinators studied by James, Roynette and Yor [3]. For more developments on \mathcal{TB} , see [4]. Part 3) of the last theorem suggests a generalization of these two notions for any parameter $a > -1$.

Notice that $\mathcal{CB}_a \subset \mathcal{CB}_b$ for every $a \leq b$, $\mathcal{TB} \subset \mathcal{SDB}$ and that $\mathcal{TB} \subset \mathcal{CB}_a \subset \mathcal{CB}$ for every $0 \leq a \leq 1$.

The simplest functions in \mathcal{CB}_a are given when taking the complete monotonic functions k of the form $k(x) = e^{-bx}$, which is the Laplace transform of δ_b , the Dirac mass at point b . Then the associated Bernstein function is

$$\phi_a(\lambda) = \int_0^\infty (1 - e^{-\lambda x}) x^{a-1} e^{-bx} dx = \begin{cases} \Gamma(a) \left(\frac{1}{b^a} - \frac{1}{(b+\lambda)^a} \right) & \text{if } a \neq 0 \\ \log(1 + \frac{\lambda}{b}) & \text{if } a = 0. \end{cases} \quad (2.3)$$

Notice that for $a \in]-1, 0[$, these functions are those associated to the so-called tempered stable processes of index $\alpha = -a$ and for $a = 0$, it is the normalized Gamma process. As settled in the next theorem, any \mathcal{CB}_a function is a conic combinations of these simple ones. This result is a straightforward consequence of theorem 2.1:

Theorem 2.3. representation of \mathcal{CB}_a -functions. *Let $a > -1$ and ϕ_a defined in (2.3). A function ϕ belongs to \mathcal{CB}_a if and only if its a Mellin convolution of ϕ_a with some measure. Namely, for $a \neq 0$*

$$\phi(\lambda) = q + d\lambda + \int_{]0, \infty[} \left(1 - \frac{t^a}{(t + \lambda)^a} \right) \nu(dt),$$

where $q, d \geq 0$, ν is a measure that integrates the function $f_a(t)$ given by (2.1). In this case, the Lévy measure associated to ϕ is $\pi(dx) = x^{a-1} \int_0^{+\infty} e^{-xt} t^a \nu(dt)$.

When $a = 0$, we have seen that we meet the well known Thorin class, then we do not recall the facts about that.

3. NEW INJECTIVE MAPPINGS FROM \mathcal{B} TO \mathcal{CB}

We recall that a \mathcal{CB} function is a Bernstein function whose Lévy measure has a density which is a completely monotonic function. We recall the connection between \mathcal{CB} functions; ϕ is a \mathcal{CB} function if and only if it admits the representation

$$\phi(\lambda) = q + a\lambda + \int_{]0, +\infty[} \frac{\lambda}{\lambda + x} \nu(dx) \quad (3.1)$$

where $q, a \geq 0$ and ν is a measure that integrates $\inf(1, 1/x)$.

Another characterization of \mathcal{CB} functions is given by theorem 6.2 of [4] and is known as the **Pick-Nevanlinna characterization of \mathcal{CB} functions** : *Let ϕ a non negative continuous function on $[0, +\infty[$ is a \mathcal{CB} function if and only if it has an analytic continuation on $\mathbf{C} \setminus]-\infty, 0]$ such that $\Im(\phi(z)) \geq 0$ for all z s.t. $\Im(z) > 0$.*

In the next theorem, we will state a representation similar to (3.1) which is valid for any function ϕ in class \mathcal{B} .

First, notice that a function ϕ in \mathcal{B} has an analytic continuation on the half plane $\{z, \Re(z) > 0\}$ which can be extended by continuity to the closed half plane $\{z, \Re(z) \geq 0\}$. We still denote ϕ this continuous continuation.

Theorem 3.1. *Let $\phi \in \mathcal{B}$, then, for all $\lambda \geq 0$,*

$$\phi(\lambda) = a\lambda + \frac{2}{\pi} \int_0^\infty \frac{\lambda}{\lambda^2 + u^2} \Re[\phi(iu)] du.$$

Proof. We suppose without loss of generality that $q = a = 0$. Recall $(C_t)_{t \geq 0}$ is a standard Cauchy process. Since $\phi(ix) = \int_{(0,\infty)} (1 - e^{-ixs}) \pi(ds)$, $x \in \mathbb{R}$, then we can write for all $\lambda > 0$:

$$\begin{aligned} \phi(\lambda) &= \int_{(0,\infty)} (1 - e^{-\lambda s}) \pi(ds) \\ &= \int_{(0,\infty)} \mathbb{E}[1 - e^{-isC_\lambda}] \pi(ds) \\ &= \int_{(0,\infty)} \left(\int_{\mathbb{R}} (1 - e^{-ius}) \frac{\lambda}{\pi(u^2 + \lambda^2)} du \right) \pi(ds) \\ &= \int_{\mathbb{R}} \frac{\lambda}{\pi(u^2 + \lambda^2)} \phi(iu) du \\ &= \int_0^\infty \frac{\lambda}{\pi(u^2 + \lambda^2)} (\phi(iu) + \phi(-iu)) du \\ &= \int_0^\infty \frac{\lambda}{\pi(u^2 + \lambda^2)} 2\Re(\phi(iu)) du \end{aligned}$$

□

Notice that $v(u) := \frac{2}{\pi} \Re(\phi(iu))$ is $[0, +\infty[$ -valued, negative definite function (see definition 4.3 and theorem 4.6 of [4]) and the identity of the preceding lemma proves that the function $\frac{v(u)}{u^2}$ is integrable at infinity. Denote by \mathcal{CB}^- the class of functions

$$\mathcal{CB}^- := \{\lambda \mapsto \psi(l) = q + d\lambda + \int_0^{+\infty} \frac{\lambda}{\lambda + u^2} v(u) du, \quad q, d \geq 0, v : [0, \infty[\rightarrow [0, \infty[\text{ negative definite function}\}$$

It is obvious that \mathcal{CB}^- is a (strict) subclass of \mathcal{CB} .

Theorem 3.2. *1) If ϕ is in \mathcal{B} , then the function $\lambda \mapsto \sqrt{\lambda}\phi(\sqrt{\lambda})$ is in \mathcal{CB}^- , more precisely,*

$$\sqrt{\lambda}\phi(\sqrt{\lambda}) = a\lambda + \int_0^\infty \frac{\lambda}{\lambda + u^2} v(u) du$$

with $v(u) = \frac{2}{\pi} \Re(\phi(iu))$.

2) Conversely, any function in \mathcal{CB}^- is of the form $\lambda \mapsto \sqrt{\lambda}\phi(\sqrt{\lambda})$ where ϕ is in \mathcal{B} .

Proof. Part 1) is given by theorem 3.1. In order to show part 2), suppose v is a (non negative) real valued function, and a negative definite function. Then, according to the Lévy-Khintchine theorem it is the Lévy Fourier exponent of an infinitely divisible distribution and it is symmetric because v is real valued. Necessarily, it is of the form $\Re(\phi(iu))$ where $\phi(iu)$ is a Lévy Fourier exponent of a spectrally

positive indefinitely divisible distribution. Now, the integrability of $\frac{v(u)}{u^2}$ at infinity implies that the quadratic component is null and that the Lévy measure integrates $x \wedge 1$ (this is easily seen by the Lévy-Khintchine formula, and can also be found in Vigon thesis [6]). Thus $\phi(iu)$ can be chosen to be the Lévy exponent of a subordinator without a drift. Then, part 1) can be applied to finish the proof. \square

Corollary 3.3. *Any Bernstein function leaves globally invariant the cônes $\{\rho e^{i\pi\alpha}; \rho \geq 0, \alpha \in [-\theta, +\theta]\}$ for every $\theta \in [0, \frac{\pi}{2}]$.*

Proof. The image of a point λ on the half line $\{\rho e^{i\pi\theta}; \rho \geq 0\}$ by the function $\lambda \mapsto \frac{\lambda}{\lambda^2 + u^2}$ (for any $u > 0$), is in the cône $\{\rho e^{i\pi\alpha}; \rho \geq 0, \alpha \in [-\theta, +\theta]\}$. This cône is convex and closed, thus, any conic combination of this function, say the integral

$$\frac{2}{\pi} \int_0^\infty \frac{\lambda}{\lambda^2 + u^2} \Re(\phi(iu)) du,$$

is in the same cône. \square

Remark 3.4. *This property has to be approached to the much stronger property fulfilled by \mathcal{CB} function : A \mathcal{CB} function has an analytic continuation on $\mathbf{C} \setminus]-\infty, 0]$ and this continuation leaves globally invariant the cônes $\{\rho e^{i\pi\alpha}; \rho \geq 0, \alpha \in [0, \theta]\}$ for any $\theta \in [0, 1[$. Moreover this property fully characterizes the \mathcal{CB} functions (the Pick-Nevenlinna characterization given above is equivalent to this property for $\theta = 1$) which is not the case for the property of \mathcal{B} - functions given in this corollary.*

In the two following results, we give some extension of theorem 3.2 by replacing the function $\lambda \mapsto \sqrt{\lambda}$ by other functions.

Corollary 3.5. *Let ϕ a \mathcal{B} -function.*

- 1) *For any function ψ such that $\lambda \mapsto \psi(\lambda^2)$ is in \mathcal{B} , the function $\lambda \mapsto \sqrt{\lambda} \phi(\psi(\lambda))$ is in \mathcal{CB}^- .*
- 2) *For any function ψ such that ψ^2 is in \mathcal{CB} , the functions and $\lambda \mapsto \psi(\lambda) \phi(\psi(\lambda))$ and $\frac{\lambda}{\psi(\lambda)} \phi(\psi(\lambda))$ are in \mathcal{CB} .*

Proof. We notice easily that we can suppose $q = a = 0$ without loss of generality and we do so.

- 1) Suppose $\lambda \mapsto \psi(\lambda^2)$ is in \mathcal{B} , then by composition, $\phi \circ \psi(\lambda^2)$ is a Bernstein function. Theorem 3.1 says that it admits the representation

$$\phi(\psi(\lambda^2)) = \int_0^\infty \frac{\lambda}{\lambda^2 + u^2} v(u) du$$

with v a completely negative function. By change of variables $\lambda \rightarrow \sqrt{\lambda}$, we then obtain :

$$\phi(\psi(\lambda)) = \int_0^\infty \frac{\sqrt{\lambda}}{\lambda + u^2} v(u) du$$

and

$$\sqrt{\lambda} \phi(\psi(\lambda)) = \int_0^{+\infty} \frac{\lambda}{\lambda + u^2} v(u) du.$$

Thus the function $\sqrt{\lambda} \phi(\psi(\lambda))$ is in \mathcal{CB}^- .

2) Compose the \mathcal{CB} function $\psi^2(\lambda)$ with the \mathcal{CB} function $\sqrt{\lambda}\phi(\sqrt{\lambda})$ and get the new \mathcal{CB} function $\psi \cdot \phi \circ \psi$ (see corollary 7.6 of [4]). This gives the first part of this assertion. Now, take the representation of ϕ of theorem 3.1, and compose with the function ψ , to get

$$\phi(\psi(\lambda)) = \int_0^\infty \frac{\psi(\lambda)}{\psi^2(\lambda) + u^2} v(u) du, \quad \left(\text{with } v(u) = \frac{2}{\pi} (\Re(\phi(iu))) \right).$$

Multiply by $\frac{\lambda}{\psi(\lambda)}$ and obtain

$$\frac{\lambda}{\psi(\lambda)} \phi(\psi(\lambda)) = \int_0^\infty \frac{\lambda}{\psi^2(\lambda) + u^2} v(u) du.$$

Since ψ^2 is in \mathcal{CB} , then, so are $\lambda \mapsto \psi^2(\lambda) + u^2$ and $\frac{\lambda}{\psi^2(\lambda) + u^2}$ is \mathcal{CB} for every u (see proposition 7.1 of [4]). Thus, the previous integral is a \mathcal{CB} -function as a conic combination of \mathcal{CB} -functions. \square

Notice that if f is a \mathcal{B} -function, then $f(\sqrt{\lambda})$ satisfies property 1). If further f is a \mathcal{CB} -function then $f(\sqrt{\lambda})$ and $\sqrt{f(\lambda)}$ both satisfy property 2).

We summarize now the properties that can be settled when composing a Bernstein function ϕ by a power function $\psi(\lambda) = \lambda^\alpha$, $\alpha \in]0, 1]$.

Proposition 3.6. *Let ϕ a function in \mathcal{B} ,*

- 1) *for all $\alpha \in]0, 1]$ the function $\lambda \mapsto \lambda^{1-\alpha} \phi(\lambda^\alpha)$ is in \mathcal{B} .*
- 2) *If $\alpha \in]0, \frac{1}{2}]$ and $\gamma \in]\alpha, 1 - \alpha]$ then $\lambda \mapsto \lambda^\gamma \phi(\lambda^\alpha)$ is in \mathcal{CB} . Moreover, if $\gamma = \frac{1}{2}$ it is in \mathcal{CB}^- .*

Proof. 1) Let $\alpha \in [0, 1[$ Denote f_α the density of a normalized α -stable distribution :

$$\int_0^{+\infty} e^{-\lambda x} f_\alpha(x) dx = e^{-\lambda^\alpha}.$$

Then, for any $t > 0$,

$$e^{-t\lambda^\alpha} = t^{-1/\alpha} \int_0^\infty e^{-\lambda x} f_\alpha(x t^{-1/\alpha}) dx. \quad (3.2)$$

Let ϕ in \mathcal{B} with Lévy-Khintchine representation:

$$\phi(\lambda) = a + b\lambda + \int_0^{+\infty} (1 - e^{-\lambda x}) \pi(dx) = a + b\lambda + \lambda \cdot \int_0^{+\infty} e^{-\lambda t} \bar{\pi}(t) dt,$$

where π is the Lévy measure and $\bar{\pi}$ its tail function $\bar{\pi}(t) = \pi([t, +\infty[)$. We have

$$\lambda^{1-\alpha} \phi(\lambda^\alpha) = a + b\lambda + \lambda \int_0^{+\infty} e^{-\lambda^\alpha t} \bar{\pi}(t) dt.$$

It is enough to prove that the integral $\int_0^\infty e^{-\lambda^\alpha t} \bar{\pi}(t) dt$ is the Laplace transform of a non increasing function. For this purpose, we use (3.2) and write

$$\int_0^\infty e^{-\lambda^\alpha t} \bar{\pi}(t) dt = \int_0^\infty t^{-1/\alpha} \left(\int_0^\infty e^{-\lambda z} f_\alpha(z t^{-1/\alpha}) dz \right) \bar{\pi}(t) dt.$$

By Fubini's theorem, we obtain that the last integral is the Laplace transform at point λ of the function

$$z \mapsto Q(z) := \int_0^\infty f_\alpha(z t^{-\frac{1}{\alpha}}) \bar{\pi}(t) t^{-\frac{1}{\alpha}} dt = \int_0^{+\infty} \bar{\pi}(z^\alpha x^{-\alpha}) f_\alpha(x) \frac{dx}{x}.$$

The second identity comes from the change of variables $t = \frac{z^\alpha}{x^\alpha}$, $x = z t^{-\frac{1}{\alpha}}$. Since the functions $z \mapsto \bar{\pi}(z^\alpha x^{-\alpha})$ are non increasing for all $x > 0$, we deduce that Q is non-increasing.

Now, for $\alpha \leq \frac{1}{2}$, the function $\psi(\lambda) = \lambda^\alpha$ satisfies the properties of corollary 3.5. Part 2) yields that the function $\lambda \mapsto \lambda^{1-\alpha} \phi(\lambda^\alpha)$ is in \mathcal{CB} , and part 1), that $\sqrt{\lambda} \phi(\lambda^\alpha)$ is in \mathcal{CB}^- .

2) If $\alpha \leq \frac{1}{2}$, then the function $\psi(\lambda) = \lambda^\alpha$ satisfy the part 1) of corollary 3.5 and we get that $\sqrt{\lambda} \phi(\lambda^\alpha)$ is in \mathcal{CB}^- .

Let now γ and α s.t. $0 < 2\alpha \leq \gamma + \alpha \leq 1$ and take the representation 3.1 of ϕ in order to obtain,

$$\lambda^\gamma \phi(\lambda^\alpha) = a \lambda^{\gamma+\alpha} + \int_0^\infty \frac{\lambda^{\gamma+\alpha}}{\lambda^{2\alpha} + u^2} v(u) du \quad v(u) = \frac{2}{\pi} \Re(\phi(iu))$$

Since $0 < 2\alpha \leq \gamma + \alpha \leq 1$, one obtains that the function $\lambda \mapsto \frac{\lambda^{\gamma+\alpha}}{\lambda^{2\alpha} + u^2}$ leaves the half plane $\{\Im(\lambda) > 0\}$ globally invariant. Thus, it is a \mathcal{CB} function for every $u > 0$. By an argument of conic combination, this property remains true for the previous integral and we can conclude that the function $\lambda \mapsto \lambda^\gamma \phi(\lambda^\alpha)$ is \mathcal{CB} .

□

4. THREE CLASSES OF RELATED BERNSTEIN FUNCTIONS

We will discuss about the classes of functions of the form $\lambda \mapsto \phi(\lambda^\alpha)$, and of the form $\lambda \mapsto \phi^\alpha(\lambda)$ and of the form $\lambda \mapsto \int_0^\infty (\frac{\lambda}{\lambda+x})^\alpha$ for any real parameter α . This part will be on line later.

REFERENCES

- [1] Bondesson, L. : Generalized Gamma Convolutions and Related Classes of Distributions and Densities. *Lect. Notes Stat.* 76, Springer-Verlag, New York (1992).
- [2] Bondesson, L. : Generalized Gamma Convolutions and Complete Monotonicity. *Probab. Theory. Rel. Fields* 85, 181-194 (1990).
- [3] James, L. F., Roynette, B., Yor, M. : Generalized Gamma Convolutions, Dirichlet means, Thorin measures, with explicit examples. *Probability Surveys* 5, 346-415 (2008).
- [4] Schilling, R.L. , Song, R. and Vondraček, Z. : *Bernstein functions*. De Gruyter, Berlin (2010).
- [5] Steutel, F.W., van Harn, K. : *Infinite divisibility of probability distributions on the real line*. Marcel Dekker, 2004.
- [6] Vigon, V. Lévy processes and Wiener-hopf factorization. PHD thesis, INSA Rouen, 2002.

SONIA FOURATI, LMI, INSA ROUEN, LPMA PARIS 6.

E-mail address: sonia.fourati@upmc.fr

WISSEM JEDIDI, DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCES OF TUNIS, CAMPUS UNIVERSITAIRE, 1060 TUNIS, TUNISIA

E-mail address: wissem.jedidi@fst.rnu.tn